

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 311/58, A61K 31/35		A1	(11) International Publication Number: WO 95/11891
			(43) International Publication Date: 4 May 1995 (04.05.95)
(21) International Application Number: PCT/SE94/01010		(74) Agent: SAMUELSSON, Britta; Astra Aktiebolag, Patent Dept., S-151 85 Södertälje (SE).	
(22) International Filing Date: 26 October 1994 (26.10.94)		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).	
(30) Priority Data: 08/144,671 28 October 1993 (28.10.93) US			
(71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): EVENDEN, John, Leslie [GB/SE]; Markvardsgatan 3, S-113 53 Stockholm (SE). HAMMARBERG, Eva, Maria [SE/SE]; Grävlingevägen 5, S-152 54 Södertälje (SE). HANSSON, Hans, Sverker [SE/SE]; Körsbärsvägen 9/1202, S-114 23 Stockholm (SE). HELLBERG, Sven, Erik [SE/SE]; Nydalavägen 60, S-152 51 Södertälje (SE). JOHANSSON, Lars, George [SE/SE]; Liljewalchsgatan 9, S-151 45 Södertälje (SE). LUND-KVIST, Johan, Rune, Michael [SE/SE]; Norrängavägen 6A, S-224 77 Lund (SE). ROSS, Svante, Bertil [SE/SE]; Hedvägen 8, S-151 52 Södertälje (SE). SOHN, Daniel, Dungan [US/SE]; Klinkervägen 11, S-151 55 Södertälje (SE). THORBERG, Seth, Olov [SE/SE]; Gullvissstigen 42, S-153 00 Järna (SE).			


Published

With international search report.

(54) Title: NOVEL (R)-5-CARBAMOYL-8-FLUORO-3-N,N-DISUBSTITUTED-AMINO-3,4-DIHYDRO-2H-1-BENZOPYRANES

(57) Abstract

Compounds of general formula (I), wherein R₁ is n-propyl or cyclobutyl; R₂ is isopropyl, tertiary butyl, cyclobutyl, cyclopentyl or cyclohexyl; R₃ is hydrogen; R₄ is hydrogen or methyl; as (R)-enantiomer and in the form of free base or pharmaceutically acceptable salts thereof, process for their preparation, pharmaceutical composition, use of and method of treatment of disorders in CNS.



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Novel (R)-5-Carbamoyl-8-fluoro-3-N,N-disubstituted-
amino-3,4-dihydro-2H-1-benzopyrans

Field of the Invention

5

The present invention relates to the new compounds,
(R)-5-carbamoyl-8-fluoro-3-N,N-disubstituted-amino-3,4-
dihydro-2H-1-benzopyrans in the form of free base or
pharmaceutically acceptable salts thereof, process for
10 their preparation, pharmaceutical compositions
containing said therapeutically active compounds and to
the use of said active compounds in therapy.

An object of the invention is to provide compounds for
15 therapeutic use, especially compounds having a highly
selective affinity for a subgroup of 5-hydroxy-
tryptamine receptors, namely the 5HT_{1A}-receptor in the
central nervous system (CNS) of mammals including man
and which compounds at the same time shows antagonist
20 activity.

It is also an object of the invention to provide
compounds with a therapeutic effect after oral
administration.

25

Prior art

Halogenated-5-subst/unsubst.carbamoyl-3-(N,N-
disubstituted-amino)-3,4-dihydro-2H-1-benzopyrans
30 intended for therapeutic use in the central nervous
system, especially having 5-HT_{1A} receptor affinity, are
already disclosed in the international patent
application WO 91/09853.

35 Background of the Invention

Various central nervous system disorders such as
depression, anxiety, etc. appear to involve the
disturbance of the neurotransmitters noradrenaline (NA)

and 5-hydroxytryptamine (5-HT), the latter also known as serotonin. The drugs most frequently used in the treatment of depression are believed to act by improving the neurotransmission of either or both of these physiological agonists. It appears that the enhancement of 5-HT neurotransmission primarily affects the depressed mood and anxiety, whereas the enhancement of noradrenaline neurotransmission affects the retardation symptoms occurring in depressed patients. The invention concerns compounds which have an effect on 5-HT neurotransmission.

Serotonin, or 5-HT, activity is thought to be involved in many different types of psychiatric disorders. For instance it is thought that an increase in 5-HT activity is associated with anxiety, while a decrease in 5-HT release has been associated with depression. Serotonin has in addition been implicated in such diverse conditions as eating disorders, gastrointestinal disorders, cardiovascular regulation and sexual behavior.

The 5-HT Receptors

The various effects of serotonin may be related to the fact that serotonergic neurons stimulate the secretion of several hormones, e.g. cortisol, prolactin, β -endorphin, vasopressin and others. The secretion of each of these other hormones appears to be regulated on a specific basis by several different 5-HT (serotonin) receptor subtypes. With the aid of molecular biology techniques, to date these receptors have been classified as 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆ and 5-HT₇ with the 5-HT₁ receptor further divided into the 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E} and 5-HT_{1F} subtypes. Each receptor subtype is involved in a different serotonin function and has different properties.

The 5-HT_{1A}-subtype Receptors

With respect to the 5-HT_{1A} subtype receptors, these receptors have also been further differentiated depending on their neuronal location thereby resulting in different modes of action and physiological functions. For example, the 5-HT_{1A} inhibitory autoreceptor is located presynaptically on cellbodies of 5-HT neurons. When 5-HT or a 5-HT_{1A} agonist activates such an inhibitory autoreceptor, the firing rate of the 5-HT neuron is depressed. Since the neuron does not fire, the release of 5-HT into the synapse is also decreased. In this case, a 5-HT_{1A} agonist acts as an artificial transmitter substance mimicking the effect of 5-HT with the result of decreasing the release of 5-HT from the central nervous system neurons. The 5-HT_{1A} agonist on the inhibitory autoreceptor thus acts as an anxiolytic or antianxiety drug.

The use of a 5-HT blocking agent at the autoreceptor will thus allow the neuron to increase its firing resulting in an increased release of 5-HT at the terminal synapses. A 5-HT_{1A} antagonist would thus improve the 5-HT neurotransmission and produce an antidepressant effect making it useful as a medication for depression.

Other localizations of 5-HT_{1A} subtype receptors also exist. These postsynaptic 5-HT_{1A} receptors are located downstream on other neurons in the synaptic region. In contrast to the inhibitory autoreceptors, activation of the postsynaptic 5-HT_{1A} receptors results in stimulation of a very characteristic behavioral syndrome in animals, the 5-HT syndrome. The agonist, partial agonist, antagonist profile of an agent on the 5-HT_{1A} postsynaptic receptor cannot be predicted from its activity profile on the 5-HT_{1A} inhibitory autoreceptor. Thus, any given 5-HT_{1A}-selective compound

may show different activity profiles on each of these receptors. For example, the D₂ antagonist Buspirone also functions as an agonist on the 5-HT_{1A} autoreceptor and acts as a weak partial agonist or antagonist on the 5-HT_{1A} postsynaptic receptor. Other compounds such as the substituted aminotetralin 8-hydroxy-2-(dipropylamino)-tetralin ("8-OH-DPAT"), which is considered the standard for 5-HT_{1A} agonist activity, is a full agonist on both the inhibitory autoreceptor and on the postsynaptic receptor.


The receptor activity profile, even within a specific subreceptor group such as 5-HT_{1A} is therefore unpredictable and may therefore result in different pharmacological profiles.

The object of the present invention is to provide compounds for therapeutic use, especially for treatment of 5-hydroxytryptamine mediated disorders in the central nervous system for instance depression, anxiety, obsessive-compulsive disorder (OCD), anorexia, bulimia, senile dementia, migraine, stroke, Alzheimer's disease, cognitive disorders, hypertension, gastrointestinal disorders, thermoregulatory and sexual disturbances, pain and for treatment of disturbances in the cardiovascular system.

Disclosure of the Invention

The object of the present invention is to provide compounds having a selective 5-HT_{1A} receptor affinity and at the same time being an antagonist, as well as having a good bioavailability.

Surprisingly, it has been found that the (R)-enantiomer of the compounds of the formula I



wherein

- 15 R_1 is n-propyl or cyclobutyl;
 R_2 is isopropyl, tertiary butyl, cyclobutyl,
cyclopentyl or cyclohexyl;
 R_3 is hydrogen;
 R_4 is hydrogen or methyl;

20 in the form of free base or pharmaceutically acceptable salts thereof, possess a high affinity to the specific subgroup of 5-HT_{1A} receptor in CNS and act as antagonists on that 5-HT_{1A} receptor, and as well show
25 sufficient bioavailability after oral administration.

Compounds of the present invention are those having the nitrogen atom on the carbamoyl group unsubstituted or mono substituted with a methyl group. The 8-fluoro
30 substituent in combination with unsubstitution or substitution with a small group as CH₃ on the carbamoyl group is very important when going from partial agonist to antagonist activity.

35 Further, the compounds of the present invention are those having at least one branched or cyclic alkyl group having 3 to 6 carbon atoms in the carbon group on the 3-amino group, thus only one of the substituents may be a normal propyl group. The branching or

cyclisation of the alkyl group on the amino group seem to be essential to obtain a sufficient bioavailability.

Furthermore, it has been found that a high affinity for
5 the 5-HT_{1A} receptor in CNS in combination with antagonist activity is strictly stereospecific to the (R)-enantiomer of the compounds of the invention.

Examples of compounds of the present invention are:

10

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-methylcarbamoyl-3,4-dihydro-2H-1-benzopyran

15

(R)-8-Fluoro-3-(N-isopropyl-N-n-propylamino)-5-carbamoyl-3,4-dihydro-2H-1-benzopyran

20

(R)-5-Carbamoyl-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

(R)-5-Carbamoyl-3-(N,N-dicyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

25

(R)-5-Carbamoyl-3-(N-cyclobutyl-N-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

(R)-5-Carbamoyl-3-(N-cyclobutyl-N-isopropylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

30

(R)-5-Carbamoyl-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

(R)-5-Carbamoyl-3-(N-cyclohexyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

35

(R)-5-Carbamoyl-3-(N-cyclopentyl-N-cyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

The compounds of the present invention are in the form of the (R)-enantiomer and in the form of free base or

pharmaceutically acceptable salt thereof. Both organic and inorganic acids can be employed to form non-toxic pharmaceutically acceptable acid addition salts of the compounds of this invention. Illustrative acids are

5 sulfuric, nitric, phosphoric, oxalic, hydrochloric, formic, hydrobromic, citric, acetic, lactic, tartaric, dibenzoyltartaric, diacetyltartaric, pamoic, ethane-


10 disulfonic, sulfamic, succinic, propionic, glycollic, malic, gluconic, pyruvic, phenylacetic, 4-aminobenzoic, anthranilic, salicylic, 4-aminosalicylic, 4-hydroxy-

15 benzoic, 3,4-dihydroxybenzoic, 3,5-dihydroxybenzoic, 3-hydroxy-2-naphtoic, nicotinic, methanesulfonic, ethanesulfonic, hydroxyethanesulfonic, benzenesulfonic, p-toluenesulfonic, sulfanilic, naphthalenesulfonic, ascorbinic, cyclohexylsulfamic, fumaric, maleic and benzoic acids. These salts are readily prepared by methods known in the art.

Methods of Preparation

20

25




30 The preparation of (R)-3-amino-5-methoxy-3,4-dihydro-2H-1-benzopyran (compound II) is described in WO93/07135. The preferred method for the introduction of fluorine is by brominating the aromatic ring, selectively, in the 8-position. Bromination can be done

35 by using bromine with or without a catalyst. Other brominating agents can be used e.g. HOBr and N-bromo amides (especially N-bromosuccinimide). Suitable solvents for bromination are acetic acid, dioxane and chlorinated solvents e.g. methylene chloride.


Before fluorination the primary amine must be fully alkylated by R_1 and by R_2 as stated above or protected by a suitable group that can be removed later e.g. dibenzyl. Introduction of the alkyl groups on nitrogen
5 can be done by reductive amination from the appropriate aldehyde or ketone using a suitable reducing agent e.g. NaCNBH_3 or catalytically with H_2 and a suitable catalyst containing palladium, platina or nickel in a suitable solvent e.g. tetrahydrofuran (THF), dioxane,
10 methanol or ethanol. Introduction of the alkyl groups can also be done by alkylation with the appropriate alkyl halide e.g. Cl, Br or I, or by an activated alcohol e.g. alkyl-mesylate or -tosylate in a suitable solvent e.g. dimethylformamide (DMF), acetone or
15 acetonitrile with a suitable base e.g. K_2CO_3 . Fluorination can then occur by lithiation of the bromo compound with an alkyllithium reagent e.g. n-butylolithium and followed by the reaction with a suitable fluorinating agent preferably a N-fluoro-N-alkyl- /arylsulfonamide e.g. N-
20 fluorobenzenesulfonimide. The solvent for this reaction can be anhydrous alkyl ethers e.g. tetrahydrofuran (THF) or diethyl ether, or non-protic solvents e.g. hexane or benzene. The temperature range can vary
25 from -100°C to room temperature but preferably -78°C to -20°C .

The compounds of the invention may be prepared from the compound (R)-3-amino-8-fluoro-5-methoxy-3,4-dihydro-2H-
30 1-benzopyran (as described above) followed by known methods such as reductive amination, N-alkylation, demethylation and conversion to the leaving group Y to obtain compound IV.

35 The compound of formula I of the invention may be prepared according to the following methods:




5 i, converting directly the compounds of formula IV



15 wherein Y is a leaving group such as trifluoromethane
 sulfonate (OSO₂CF₃), halide e.g. Cl, Br or I by a
 catalytic cycle using a zerovalent transition metal (M)
 such as Pd or Ni, which may be generated in situ and
 undergoes an oxidative addition to the aryl-Y-bond.
 Treatment with carbon monoxide followed by amination
 20 with the proper amine (ammonia or methylamine) give the
 compounds of formula I, whereafter if desired it is
 converted to a salt.

ii, Alternatively, the compounds of formula IV is converted to the compounds of formula V



wherein Z is Cl, Br, OH or OR_p, where R_p is C₁-C₆ alkyl, by a catalytic cycle using a zerovalent transition metal, with ability to undergo oxidative addition to aryl-Y-bonds e.g. the aryl-OSO₂CF₃ bonds. The aryl-CO-metal-Y complex is formed by treatment with carbon monoxide (CO).

Further reagents are an alcohol such as methanol, ethanol, a tertiary amine such as a trialkylamine e.g. triethylamine in an inert organic solvent preferentially a polar aprotic solvent such as dimethylformamide (DMF), dimethylsulfoxide (DMSO), dioxane, tetrahydrofuran (THF), acetone, acetonitrile etc. The reaction is normally performed at a temperature between +40 to +120°C and at a pressure between 100 to 500 kPa (ii). This is optionally followed by hydrolysis and treatment with a thionyl halide e.g. thionyl chloride, to obtain the corresponding acid halide derivative.

The compounds of formula V is aminated (ii_b) with the proper amine (ammonia or methylamine) in a solvent e.g. toluene, methylene chloride, benzene, water at reflux temperature or between 0-100°C to give the compounds of formula I.

Pharmaceutical preparations

5 According to the present invention the compound of the
invention will normally be administered orally,
rectally or by injection, in the form of pharmaceutical
preparations comprising the active ingredient either as
a free base or a pharmaceutically acceptable non-toxic
acid addition salt, e.g. the hydrochloride,
10 hydrobromide, lactate, acetate, phosphate, sulfate,
sulfamate, citrate, tartrate, oxalate and the like in a
pharmaceutically acceptable dosage form. The dosage
form may be a solid, semisolid or liquid preparation.
Usually the active substance will constitute between
15 0.1 and 99% by weight of the preparation, more
specifically between 0.5 and 20% by weight for
preparations intended for injection and between 0.2 and
50% by weight for preparations suitable for oral
administration.

20 To produce pharmaceutical preparations containing the
compound of the invention in the form of dosage units
for oral application, the selected compound may be
mixed with a solid excipient, e.g. lactose, saccharose,
25 sorbitol, mannitol, starches such as potato starch,
corn starch or amylopectin, cellulose derivatives, a
binder such as gelatine or poly-vinylpyrrolidone, and a
lubricant such as magnesium stearate, calcium stearate,
polyethylene glycol, waxes, paraffin, and the like, and
30 then compressed into tablets. If coated tablets are
required, the cores, prepared as described above, may
be coated with a concentrated sugar solution which may
contain e.g. gum arabic, gelatine, talcum, titanium
dioxide, and the like. Alternatively, the tablet can be
35 coated with a polymer known to the man skilled in the
art, dissolved in a readily volatile organic solvent or
mixture of organic solvents. Dyestuffs may be added to
these coatings in order to readily distinguish between
tablets containing different active substances or

different amounts of the active compound.

For the preparation of soft gelatine capsules, the active substance may be admixed with e.g. a vegetable
5 oil or poly-ethylene glycol. Hard gelatine capsules may contain granules of the active substance using either the above mentioned excipients for tablets e.g. lactose, saccharose, sorbitol, mannitol, starches (e.g. potato starch, corn starch or amylopectin), cellulose
10 derivatives or gelatine. Also liquids or semisolids of the drug can be filled into hard gelatine capsules.

Dosage units for rectal application can be solutions or suspensions or can be prepared in the form of supposi-
15 tories comprising the active substance in a mixture with a neutral fatty base, or gelatine rectal capsules comprising the active substance in admixture with vegetable oil or paraffin oil. Liquid preparations for oral application may be in the form of syrups or
20 suspensions, for example solutions containing from about 0.2% to about 20% by weight of the active substance herein described, the balance being sugar and mixture of ethanol, water, glycerol and propylene glycol. Optionally such liquid preparations may contain
25 colouring agents, flavouring agents, saccharine and carboxymethyl-cellulose as a thickening agent or other excipients known to the man in the art.

Solutions for parenteral applications by injection can
30 be prepared in an aqueous solution of a water-soluble pharmaceutically acceptable salt of the active substance, preferably in a concentration of from about 0.5% to about 10% by weight. These solutions may also contain stabilizing agents and/or buffering agents and
35 may conveniently be provided in various dosage unit ampoules.

Suitable daily doses of the compound of the invention in therapeutical treatment of humans are about 0.01-100 mg/kg bodyweight at peroral administration and 0.001-

100 mg/kg bodyweight at parenteral administration.

The invention is illustrated by the following working examples.

5

Preparation of Starting Materials

Preparation 1

10 (R)-3-Amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

a) (R)-3-Amino-8-bromo-5-methoxy-3,4-dihydro-2H-1-benzopyran

15 (R)-3-Amino-5-methoxy-3,4-dihydro-2H-1-benzopyran
(25 g, 0.14 mol) and anhydrous sodium acetate (34 g,
0.42 mol) were dissolved in acetic acid (500 mL).
Bromine (23.4 g, 0.15 mol), dissolved in acetic acid
(500 mL), was added dropwise at room temperature. The
20 addition of bromine continued for about 7 days. The
solvent was removed in vacuo and the residue was
dissolved in sodium hydroxide (25%)/diethyl ether
(exothermic reaction, the mixture was cooled in an ice-
bath). The layers were separated and the alkaline
25 water-phase was extracted thrice with diethyl ether.
The combined ether layers were dried (Na_2SO_4) and the
solvent was removed in vacuo to give 35.5 g of an oily
residue. The oil was dissolved in diethyl ether and the
solution was cooled with an ice-bath (0°C). HCl,
30 dissolved in diethyl ether, was added dropwise until
the suspension became acidic (controlled with pH
paper). The crystals were filtered and then
recrystallized from methanol to give the title compound
in 70% yield (28.5 g). Mp: 281-282°C. The HCl salt was
35 partitioned between diethyl ether and 2 M NH_3 (aq) and
the free base was obtained by extractions of the
alkaline water phase with diethyl ether. $[\alpha]_D^{21} = +40^\circ$
(C=0.1, HCl salt in MeOH). GC-MS (70eV) M+1 = 259
(53%).

b) (R)-8-Bromo-3-(N,N-dibenzylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran

5 (R)-3-Amino-8-bromo-5-methoxy-3,4-dihydro-2H-1-benzopyran (11.5 g, 44 mmol) was dissolved in 400 mL anhydrous acetonitrile and to the reaction were benzyl bromide (13 mL, 110 mmol), anhydrous potassium carbonate (grounded) (16 g, 116 mmol), and a catalytic
10 amount of KI added and then heated to 85°C for 48 h. The solvent was removed in vacuo, the remains were taken into a 2 M solution of NH₃ and then extracted twice with ether. The combined ether portions were treated with brine, dried (MgSO₄), filtered, and the
15 solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: CH₂Cl₂) gave 15 g (78% yield) of the title compound as a clear oil. $[\alpha]_D^{21} = -15.5^\circ$ (C=0.1, CHCl₃). GC-MS (70eV) M=437 (15%).

20

c) (R)-3-(N,N-Dibenzylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-8-Bromo-3-(N,N-dibenzylamino)-5-methoxy-3,4-dihydro-2 H-1-benzopyran (4.35 g, 9.9 mmol) was
25 dissolved in 45 mL of anhydrous THF and cooled to -78°C. To this was a 1.6 M n-BuLi solution (6.8 mL, 10.9 mmol) added dropwise and allowed to stir at -78°C for 1 h. N-Fluorobenzenesulfonimide (3.8 g, 11.9 mmol),
30 dissolved in 30 mL anhydrous THF, was added dropwise under 45 min and allowed to stir at -78°C for 1 h. The reaction was stopped by adding 3 mL saturated NH₄Cl followed by 9 mL of a solution comprised of 2 g of NH₂OH·HCl and 8 g of Na₂CO₃ in 100 mL of H₂O and
35 allowing the reaction to warm to room temperature. A 2 M NH₃ solution was added and the reaction was extracted twice with ether, treated with brine, dried (Na₂SO₄), filtered, and evaporated in vacuo to give the crude product.

Purification of the 8-fluoro (desired) from the 8-hydrogen (15 %) compound was carried out by a crude chromatography (eluent: 25 % CH₂Cl₂/hexane) to give 1.78 g. The crude was rechromatographed on silica
5 (eluent: 3% acetone/hexane) to give 1.50 g (40% yield) of the title compound as a clear oil. $[\alpha]_D^{21} = -112.1^\circ$ (C=0.1, CHCl₃). GC-MS (70 eV) M=377 (37%).

10 **d) (R)-3-Amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran**

(R)-3-(N,N-Dibenzylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (13.0 g, 34.4 mmol) was dissolved in 265 mL methanol and 115 mL THF. To this
15 was 10 % Pd/C (4 g) and ammonium formate (51.5 g, 0.817 mol) added. The reaction was heated to 50°C for 2.5 h. The reaction was filtered and the solvent was removed in vacuo, the remains were taken into a 2 M solution of NaOH and then extracted twice with ether. The combined
20 ether portions were treated with brine, dried (Na₂SO₄), filtered, and the solvent removed in vacuo to give 6.2 g (91 % yield) of the title compound as a clear oil. $[\alpha]_D^{21} = -15.3^\circ$ (C=1, CHCl₃). GC-MS (70 eV) M=197 (51%).

25

Preparation 2

(R)-8-Fluoro-3-(N-isopropyl-N-propylamino)-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran

30 **a) (R)-8-Bromo-3-(N-isopropylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran**

(R)-3-(N-Isopropylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran (4.02 g, 18.2 mmol) and anhydrous sodium acetate were dissolved in acetic acid (80 mL). To the
35 stirred mixture was bromine (0.93 mL, 18.2 mmol) dissolved in acetic acid (40 mL) added dropwise under 1.5 h.

The solvent was removed in vacuo, taken into a 2 M NaOH

solution and extracted twice with diethyl ether. The combined ether portions were treated with brine, dried (Na_2SO_4), filtered, and the solvent was removed in vacuo to give the crude residue. The hydrochloride salt was made by dissolving the pure base in diethyl ether and adding an excess of an ethereal HCl solution to give a white solid. The salt was recrystallized twice from ethanol/diethyl ether to give 3.8 g (62% yield). Mp: 257-8°C. $[\alpha]^{21}_{\text{D}} = -97.7^\circ$ (C=0.1, CHCl_3). The free base was made from the hydrochloride salt to give an oil. GC-MS (70 eV) M=301 (100%).

b) (R)-8-Bromo-3-(N-isopropyl-N-propylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-8-Bromo-3-(N-isopropylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran (3.8 g, 11.3 mmol) was dissolved in anhydrous methanol (80 mL) and to this was propanal (8.1 mL, 0.113 mol) added. The reaction was cooled (ice-bath) then sodium cyanoborohydride (1.3 g, 20.3 mmol) was added, the pH was adjusted to 5, and the reaction was allowed to stir at room temperature overnight.

The solvent was removed in vacuo, the remains were taken into a 1 M solution of Na_2CO_3 and then extracted twice with diethyl ether. The combined ether portions were treated with brine, dried (Na_2SO_4), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 7% ethyl acetate/hexane) gave 3.75 g (97% yield) of the title compound as a clear oil. $[\alpha]^{21}_{\text{D}} = -82.5^\circ$ (C=0.1, CHCl_3). GC-MS (70 eV) M=343 (29%).

The hydrochloride salt was made by dissolving the base in diethyl ether and dropping an excess of an ethereal HCl solution. The salt was recrystallized from ethanol/diethyl ether to give a white solid. Mp: 177-9°C.

c) (R)-8-Fluoro-3-(N-isopropyl-N-propylamino)-5-

methoxy-3,4-dihydro-2H-1-benzopyran

(R)-8-Bromo-3-(N-isopropyl-N-propylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran (2.3 g, 6.72 mmol) was dissolved in anhydrous THF (25 mL) and cooled to -78°C. To this was a 1.6 M n-BuLi solution (4.83 mL, 7.73 mmol) added dropwise and allowed to stir at -78°C for 1 h. N-Fluorobenzenesulfonimide (2.55 g, 8.06 mmol), dissolved in anhydrous THF (15 mL), was added dropwise under 20-30 min and allowed to stir at -78°C for 4 h. The reaction was stopped by adding 1 mL of a saturated aqueous NH_4Cl solution followed by 3 mL of a solution comprised of 2 g of $\text{NH}_2\text{OH}\cdot\text{HCl}$ and 8 g of Na_2CO_3 in 100 mL of H_2O and allowing the reaction to warm to room temperature. A 2 M NH_3 solution was added and the reaction was extracted, twice, with diethyl ether, treated with brine, dried (Na_2SO_4), filtered, and evaporated in vacuo to give the crude product. Chromatography on silica (eluent: chloroform) gave 1.0 g (53% yield) of the title compound as a clear oil. $[\alpha]_D^{21} = -89.2^\circ$ ($C=0.1$, CHCl_3). GC-MS (70 eV) $M=281$ (32%).

The hydrochloride salt was made by dissolving the pure base in diethyl ether and dropping an excess of an ethereal HCl solution to give a white solid (sinters at 80°C).

d) (R)-8-Fluoro-3-(N-isopropyl-N-propylamino)-5-hydroxy-3,4-dihydro-2H-1-benzopyran

(R)-8-Fluoro-3-(N-isopropyl-N-propylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran hydrochloride (1.03 g, 3.24 mmol) was dissolved in anhydrous CH_2Cl_2 (30 mL) and cooled to -40°C. To the solution was BBr_3 (0.77 mL, 8.1 mmol), dissolved in anhydrous CH_2Cl_2 (5 mL), added dropwise. The cooling-bath was removed and after 3 h at room temperature the reaction was complete. The reaction was poured out onto an ice/2 M NH_3 solution and the mixture was extracted, twice, with

diethyl ether. The combined ether portions were treated with brine, dried (Na_2SO_4), filtered, and the solvent removed in vacuo to give the crude residue.

Chromatography on silica (eluent: 20% ethyl

5 acetate/hexane) gave 0.84 g (97% yield) of the title compound as a clear oil. $[\alpha]_D^{21} = -94.2^\circ$ ($C=0.1$, CHCl_3). GC-MS (70 eV) $M=267$ (26%).

The hydrochloride salt was made by dissolving the pure base in diethyl ether and dropping an excess of an
10 ethereal HCl solution. The salt was recrystallized from CHCl_3 /diethyl ether/ethyl acetate to give a white solid. Mp: $220-2^\circ\text{C}$.

e) (R)-8-Fluoro-3-(N-isopropyl-N-propylamino)-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran
15

(R)-8-Fluoro-3-(N-isopropyl-N-propylamino)-5-hydroxy-3,4-dihydro-2H-1-benzopyran (0.71 g, 2.66 mmol) and collidine (0.49 mL, 3.72 mmol) were dissolved in
20 anhydrous CH_2Cl_2 (25 mL) and cooled to -40°C . Trifluoromethanesulfonic anhydride (0.54 mL, 3.2 mmol) was added dropwise and allowed to warm to ambient temperature, and after coming to 0°C the reaction was done. The reaction was diluted with CH_2Cl_2 and washed
25 with a saturated aqueous NaHCO_3 solution, dried (MgSO_4), filtered, and evaporated in vacuo to give a crude residue. Chromatography on silica (eluent: CH_2Cl_2) gave 0.82 g (77% yield) of the title compound as a clear oil. $[\alpha]_D^{21} = -85.5^\circ$ ($C=0.1$, CHCl_3). GC-MS
30 (70 eV) $M=399$ (6%).

Example 1

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-N-methylcarbamoyl-3,4-dihydro-2H-1-benzopyran
35

a) (R)-3-N-Cyclopentylamino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-3-Amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (1.5 g, 7.6 mmol), acetic acid (0.45 g, 7.6 mmol), and cyclopentanone (2.5 g, 3 mmol) were dissolved in 30 ml of methanol. With stirring, sodiumcyanoborohydride (0.8 g, 13 mmol) was added in portions under a few minutes. Stirring was continued for 2 hours. A GC sample showed 100% of a new product. The solvent was evaporated and water, 2 M NH₃ and EtOAc were added. The organic layer was separated and washed with water. The layer was dried with Na₂SO₄ and evaporated to give 1.3 g (64% yield) of a colourless oil. GC/MS with the molecular peak of 265 confirmed the title compound.

b) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-3-N-Cyclopentylamino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (1.3 g, 5 mmol), acetic acid (0.3 g, 5 mmol) and propionaldehyde (1.5 g, 25 mmol) were dissolved in 30 ml of methanol. With stirring, sodiumcyanoborohydride (0.8 g, 13 mmol) was added in portions under a few minutes and stirring was continued. After 3 hours a GC sample showed 100% of a new product. The solvent was evaporated and water, 2 molar NH₃ and EtOAc were added. The organic layer was separated and washed neutral with water. The layer was dried with Na₂SO₄ and evaporated to give 1 g (65% yield) of a colourless oil. GC/MS with the molecular peak of 307 confirmed the title compound.

c) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (1 g, 5 mmol) was dissolved in 25 ml of CH₂Cl₂. An excess of etheric HCl was added to form the HCl salt. A solution of BBr₃ (4 g, 15 mmol) in 10 ml of CH₂Cl₂ was prepared and added

dropwise under 10 minutes with stirring (ice-bath). The reaction mixture was allowed to reach room temperature during continued stirring for 6 hours and the mixture was poured out into ice water and made alkaline by adding ammonia. The organic layer was separated, dried with Na_2SO_4 and evaporated to afford a dark brown oil. Chromatography (SiO_2 , di-isopropyl ether and hexane 1+1) afforded 1.1 g of an colourless oil. The HCl salt was prepared from the base and etheric HCl and rerystallized from acetonitrile to give 0.85 g (52% yield). Mp 220-221°C.

d) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran (0.7 g, 3 mmol) was dissolved in 25 ml of CH_2Cl_2 and triethylamine (0.3 g, 3 mmol) was added. The solution of trifluoromethanesulfonic anhydride (1 g, 4 mmol) in 5 ml of CH_2Cl_2 was added dropwise under 10 min (-20°C). Stirring was continued for 1 hour. The reaction mixture was poured out into ice water and the pH was adjusted to 8 by addition of ammonia and extracted by ether. The organic layer was separated, dried with Na_2SO_4 and evaporated to afford a brown oil. Chromatography (SiO_2 , CH_2Cl_2 + hexane, 1 + 3) afforded 0.5 g (44% yield) of a colourless oil. GC/MS with the molecular peak of 425 confirmed the title compound.

e) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-N-methylcarbamoyl-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (0.5 g, 1 mmol) was dissolved in 15 ml of 1,4-dioxane. Palladium II acetate (10 mg), 1,3-bis(diphenylphosphino)-propane (20 mg), and methylamine (0.15 g, 5 mmol) were added and the mixture was stirred

in carbon monoxide atmosphere over night at 70°C. Evaporation and chromatography (SiO₂, diethyl ether + hexane 1+3) afforded the final compound as a colourless oil. The HCl salt was prepared to give 0.24 g (65% yield) of white crystals. Mp 108°C.

Example 2

(R)-8-Fluoro-3-(N-isopropyl-N-n-propylamino)-5-carbamoyl-3,4-dihydro-2H-1-benzopyran

10

a) Methyl (R)-8-fluoro-3-(N-isopropyl-N-n-propylamino)-3,4-dihydro-2H-1-benzopyran- 5-carboxylate

(R)-8-Fluoro-3-(N-isopropyl-N-n-propylamino)-5-trifluoromethanesulfonyloxy-3,4-dihydro-2H-1-benzopyran (2.4 g, 6.0 mmol), triethylamine (1.3 g, 12.9 mmol), 1,3-bis(diphenylphosphino)propane (95 mg, catalytic amount), palladium(II)acetate (48 mg, catalytic amount) and DMF/MeOH (30 mL, 3:1) were mixed in a 50 mL three necked round bottom flask. The flask was evacuated followed by the inlet of CO (repeated two times). The reaction mixture was stirred at 70°C for 7.5 hours. The solvent was removed in vacuo and the residue was dissolved in diethyl ether/sat. NaHCO₃. The layers were separated and the water phase was extracted once with ether. The combined ether layers were dried (MgSO₄) and the solvent was removed in vacuo to give a crude which was purified by flash chromatography (SiO₂, hexane/EtOAc, 9:1) to give 1.3 g of the title compound (71% yield).

b) (R)-8-Fluoro-3-(N-isopropyl-N-n-propylamino)-5-carbamoyl-3,4-dihydro-2H-1-benzopyran

Methyl (R)-8-Fluoro-3-(N-isopropyl-N-n-propylamino)-3,4-dihydro-2H-1-benzopyran-5-carboxylate (1.3 g, 4.2 mmol) and KOH (0.52 g, 8.4 mmol) were mixed in methanol (6 mL) and refluxed for 2.5 hours. The solvent was removed in vacuo. The residue was dissolved in water

and made acidic by the addition of 2M HCl. The solvent was removed in vacuo. The residue was dissolved in SOCl₂ (30 mL) and refluxed for 2.5 hours. The solvent was removed in vacuo. The residue was dissolved in CH₂Cl₂ and the solvent was removed in vacuo again (repeated three times in order to remove the excess of SOCl₂). The residue was then dissolved in diethyl ether (50 mL). The solution was cooled to -30°C before NH₃ (g) was bubbled through it. Water was added, the layers were separated and the water-phase was extracted with ether. The combined ether layers were dried (K₂CO₃) and the solvent was removed in vacuo to give a crude which was purified by flash chromatography (SiO₂, EtOAc/hexane, 1:1) to give 1.0 g of the title compound (yield 80%). Recrystallization from EtOAc/hexane gave crystals with Mp: 139-140°C.

Example 3

(R)-3-(N-tert-Butyl-N-n-propylamino)-5-carbamoyl-8-fluoro-3,4-dihydro-2H-1-benzopyran

a) (R)-8-Fluoro-5-methoxy-3-[N-(4-methoxybenzylidene)-amino]-3,4-dihydro-2H-1-benzopyran

(R)-3-Amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (7.85 g, 39.8 mmol), 4-methoxybenzaldehyde (5.42 g, 39.8 mmol), anhydrous potassium carbonate (10.1 g) and absolute EtOH (200 mL) were stirred over night at reflux temperature. The solvent was evaporated in vacuo and ether (500 mL) was added. After stirring for 15 min the salt was filtered off and the solution was concentrated in vacuo to give an off-white solid (12.4 g). Recrystallization from i-Pr₂O- hexane gave 10.8 g (86% yield) of the title compound as colourless needles. Mp: 96.8-97.3°C.
[α]_D²¹ = +20.1° (C=1; CHCl₃). GC-MS (70 eV) M= 315 (58%).

b) (R)-8-Fluoro-3-hydroxylamino-5-methoxy-3,4-dihydro-2H-1-benzopyran

3-Chloroperoxybenzoic acid (85%; 7.6 g, 37.6 mmol) was
5 added in portions to a stirred and cooled solution
(+ 4°C) of (R)-8-fluoro-5-methoxy-3-[N-(4-
methoxybenzylidene)-amino]-3,4-dihydro-2H-1-benzopyran
(10.8 g, 34 mmol) and methylene chloride (65 mL). The
mixture was stirred over night at room temperature.
10 Precipitated 3-chlorobenzoic acid was filtered off and
the clear yellow filtrate was concentrated in vacuo.
The oily residue was taken up in a solution of
hydroxylamine hydrochloride (2.83 g, 40.8 mmol) and
anhydrous methanol (60 mL) and the resulting solution
15 was stirred at room temperature for 2 h. The solvent
was evaporated to give a thick orange oil. Water was
added to the oil, pH was adjusted to 8-9 with saturated
aqueous Na₂CO₃ and the mixture was washed with ether
(3x150 mL). The organic phases were combined, washed
20 with brine, dried (Na₂SO₄), filtered and concentrated
in vacuo. The crude product was flash chromatographed
on silica with EtOAc (15 to 50%) in hexane as eluent.
The resulting impure product was flash chromatographed
a second time on silica with EtOH-CHCl₃ (1:99) as
25 eluent to give 6.45 g (89% yield) of the title compound
as a colourless crystalline solid. Mp: 111-113°C.
[α]_D²¹ = +66.4° (C= 1.3; CHCl₃). GC-MS (70 eV) M= 213
(56%).

30 c) (R)-3-tert-Butylamino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-8-Fluoro-3-hydroxylamino-5-methoxy-3,4-dihydro-2H-
1-benzopyran (6.30 g, 29.6 mmol), anhydrous sodium
35 sulfate (20 g) and acetone (500 mL) were refluxed under
nitrogen for 4 days until TLC indicated a complete
reaction. The salt was filtered off, ether (300 mL)
was added to the filtrate and the solution, still
containing finely suspended salt, was filtered through

a sintered glass filter (grade 4). The clear filtrate was concentrated in vacuo. Dry (sieves 3 Å) benzene (50 mL) was added and the resulting solution was concentrated in vacuo (finally on the pump). The glassy residue was dissolved in dry (sieves 3 Å) benzene (150 mL) under nitrogen and the solution was cooled on an ice-bath (+4°C). MeMgBr in Et₂O (3.0 M; 32.0 mL, 96 mmol) was added to the stirred solution above at a rate that kept the internal temperature below +5°C (the reaction is exothermic). After the addition was complete (30 min) the solution was stirred at +4°C for 0.5 h. The cooling bath was taken away and 15 min later the solution was poured on saturated NaHCO₃ and ice (300 mL total). The mixture was washed repeatedly with ether (3x150 mL). The organic phases were combined, washed with brine, dried (Na₂SO₄), filtered and concentrated in vacuo. Flash chromatography on silica [eluent: EtOAc (2 and 10%) in CHCl₃] gave 2.9 g. of the tert -butyl hydroxylamine derivative. The latter was dissolved in CS₂ (100 mL) under nitrogen and the solution was stirred at room temperature for 4.5 h. The solvent was evaporated in vacuo to give an orange oil. Acetone (approx. 50 mL) was added and the solution was stirred for a short time (15 min) at room temperature (to precipitate elemental sulfur), then filtered and concentrated to give an oil. Flash chromatography on silica [eluent: EtOAc (10 to 25%) in hexane] gave 2.34 g (31% total yield) of the title compound as a yellow oil.

[α]_D²¹ = -82.8° (C = 1; CHCl₃). GC-MS (70 eV) M = 253 (53%).

d) (R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-3-tert-Butylamino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (2.20 g, 8.7 mmol), allyl bromide (7.5 mL, 87 mmol), finely ground anhydrous potassium carbonate (6.0 g, 43 mmol) and dry DMF (6.0 mL) were

stirred under nitrogen at 65°C. After 70 h GC-analysis indicated partial conversion of the starting material (67% product vs. 30% starting material). At this stage the reaction was interrupted. The salt was sucked off, washed with a small portions of DMF and the clear filtrate was concentrated. The oil thus obtained was partitioned between saturated aqueous Na₂CO₃ and diethyl ether (4x70 mL). The organic phases were combined, washed with brine, dried (MgSO₄), filtered and concentrated in vacuo. Flash chromatography on silica [eluent: acetone (2 and 15%) in hexane] gave 0.80 g of starting material and 1.47 g (87% yield based on recovered starting material) of the allylated product as a colourless oil. $[\alpha]_D^{21} = -77.6^\circ$ (C= 0.8, CHCl₃). GC-MS (70 eV) M= 293 (40%).

The allylated product (1.30 g) was mixed with DMF (50 mL) and 5% Rh on alumina (0.090 g) and hydrogenated at ambient pressure and temperature (21°C). After 5 h the reaction was complete according to GC and TLC. The catalyst was filtered off on Celite, the pad was washed with small portions of DMF and the filtrate was concentrated in vacuo. Flash chromatography on silica of the crude product [eluent: EtOAc (0 and 3%) in CH₂Cl₂] gave 1.27 g (97% yield) of the saturated compound.

GC-MS (70 eV) M= 295 (28%). $[\alpha]_D^{21}$ (base) = -83.4° (C= 0.9; CHCl₃).

The base was dissolved in dry diethyl ether, the solution was cooled on an ice bath and an excess of ethereal HCl was added to the stirred solution. The salt was filtered off, washed with dry diethyl ether and dried in vacuo at 50°C to give 1.39 g (98% yield) of the title compound, as white crystals.

Mp: 175-176°C.

e) (R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran hydrochloride (1.3 g, 3.9 mmol) in dry methylene chloride (40 mL) under nitrogen was cooled on a dry ice-EtOH bath to -50°C. Boron tribromide (0.75 mL, 7.8 mmol) was added dropwise (in 1 min) to the stirred solution above. Five minutes after the addition of boron tribromide was complete, the dry-ice bath was changed to an ice bath (+4°C). After stirring for 4 h at the same temperature the solution was poured on ice (100 g) and solid NaHCO₃ was added to adjust pH to 8-9. When the ice had melted the mixture was extracted with ether (4x75 mL). The ether extracts were combined, washed with brine, dried (Na₂SO₄), filtered and concentrated in vacuo to afford 1.1 g (96% yield) of the title compound as a pale yellow oil.

[α]_D²¹ = -91.7° (C= 1.0; CHCl₃). GC-MS (70 eV) M= 281 (6%).

f) (R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran (1.0 g, 3.6 mmol) and 2,4,6-collidine (0.52 mL, 3.9 mmol) were dissolved in anhydrous methylene chloride (40 mL) and cooled to -30°C. Trifluoromethanesulfonic anhydride (0.66 mL, 3.9 mmol) dissolved in anhydrous methylene chloride (10 mL) was added dropwise during 20 min. The solution was allowed to warm to ambient temperature and after coming to 0°C the reaction was done. The reaction was diluted with methylene chloride and washed with saturated aqueous NaHCO₃ (50 mL). The aqueous phase was re-extracted with ether (2x40 mL). The combined organic phases were dried (MgSO₄), filtered, and concentrated in vacuo to give a crude residue. Flash chromatography

on silica [eluent: EtOAc-hexane (3:97)] gave 1.40 g (95% yield) of the title compound as a colourless oil. $[\alpha]_D^{21} = -73.7^\circ$ (C= 1.1; CHCl₃). GC-MS (70 eV) M= 413 (1%).

5

g) Methyl (R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate

(R)-3-(N-tert-Butyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (1.4 g, 3.3 mmol) and triethylamine (1.0 mL, 7.4 mmol) were dissolved in a solution of DMF/MeOH (6:2; 30 mL) and then degassed followed by the inlet of carbon monoxide (4x). With a slight positive pressure of carbon monoxide, palladium(II)-acetate (0.030 g) and 1,3-bis(diphenylphosphino)propane (0.060 g) were added and the reaction mixture was degassed and subjected to carbon monoxide once again. The reaction was heated to 70°C (oil-bath temperature) under carbon monoxide atmosphere with vigorous stirring for 12 h. GC indicating an incomplete reaction (68% product vs. 21% starting material), the solution was cooled and then filtered through Celite. More palladium(II)-acetate (0.030 g) and 1,3-bis(diphenylphosphino)propane (0.060 g) were added and the reaction was resumed as described above. After another 3 h GC indicated a slight improvement of the ratio (77% vs. 12%) and the reaction was allowed to cool. The following day the solvent was removed in vacuo. The remaining red-brown oil was taken into saturated aqueous NaHCO₃ and then extracted with EtOAc (3x50 mL). The combined organic phases were washed with brine, dried (MgSO₄), filtered and concentrated in vacuo to give the crude ester. Flash chromatography on silica [eluent: EtOAc (15 and 30%) in hexane] gave 0.178 g of starting material and 0.842 g (89% yield based on recovered starting material) of the title compound as a clear oil. $[\alpha]_D^{21} = -121.1^\circ$ (C= 0.9; CHCl₃). GC-MS (70 eV) M= 323 (14%).

h) (R)-3-(N-tert-Butyl-N-n-propylamino)-5-carbamoyl-8-fluoro-3,4-dihydro-2H-1-benzopyran

Methyl (R)-3-(N-tert-butyl-N-n-propylamino)-8-fluoro-
5 3,4-dihydro-2H-1-benzopyran-5-carboxylate (0.84 g, 2.6 mmol), methanol (10 mL) and 1.7 M aqueous NaOH (3.0 mL, 5.2 mmol) were refluxed for 3 h. The clear solution was cooled, the methanol was stripped, the aqueous remains were washed twice with ether-hexane (1:1), then
10 acidified with 2 M HCl (pH \leq 2). The water was evaporated in vacuo and the remaining salt was dried in vacuum at 50°C for 2 h. Dry methylene chloride (20 mL) and thionyl chloride (3.0 mL, 41 mmol) were added, the mixture refluxed under nitrogen for 11 h. The
15 volatiles were evaporated, more dry methylene chloride was added and evaporated. This was repeated once. The crude acid chloride was dissolved (suspended) in dry methylene chloride (50 mL) and added dropwise to stirred concentrated aqueous ammonia (40 mL) cooled on
20 an ice bath. The mixture was allowed to warm to ambient temperature, the organic phase was separated, and the aqueous phase was washed with methylene chloride (100 mL) and ether (50 mL). The organic portions were combined, dried (MgSO₄), filtered and concentrated to
25 give the crude amide. Flash chromatography on silica [eluent: EtOAc-hexane (4:5)] gave 0.73 g (91% yield) of (R)-3-(N-tert-butyl-N-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxamide as a solid.
Mp: 70-75°C.
30 $[\alpha]_D^{21} = -132.4^\circ$ (C= 1.0; CHCl₃). GC-MS (70 eV) M= 308 (3%).

The base was dissolved in dry ether, the solution was cooled on a dry-ice bath (-20°C) and an excess of
35 ethereal HCl was added to the stirred solution. The salt was filtered off, washed with dry ether and dried in vacuo at 50°C to give 0.78 g (96% yield) of the hydrochloride salt as white crystals.
Mp: 120°C sinters.

Example 4(R)-5-Carbamoyl-3-(N,N-dicyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

- 5 a) (R)-3-(N-Cyclobutylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran and
(R)-3-(N,N-Dicyclobutylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran
- 10 (R)-3-Amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (1.67 g, 8.47 mmol) was dissolved in anhydrous methanol (20 mL) and to this was cyclobutanone (5.0 g, 71.3 mmol) added. The reaction was cooled (ice-bath) then sodium cyanoborohydride
- 15 (0.96 g, 15.3 mmol) was added and the reaction was allowed to stir at room temperature overnight. After 24 h the pH was adjusted to 4-5 with acetic acid and the reaction was allowed to stir for one more day. The solvent was removed in vacuo, the remains were
- 20 taken into a 2 M solution of NH_3 and then extracted thrice with diethyl ether. The combined ether portions were dried (Na_2SO_4), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 15% ethyl acetate/hexane for
- 25 dialkylated product followed by ethyl acetate for monoalkylated product) gave 1.01 g (48% yield) of the monoalkylated title compound as a clear oil [GC-MS (70 eV) $M = 251$ (6%)]. and 0.71 g (27% yield) of the dialkylated title compound as a clear oil.
- 30 $[\alpha]_D^{21} = -101.0^\circ$ ($C = 0.1$; CHCl_3). GC-MS (70 eV) $M = 305$ (3%).
- b) (R)-8-Fluoro-3-(N,N-dicyclobutylamino)-5-hydroxy-3,4-dihydro-2H-1-benzopyran
- 35 (R)-8-Fluoro-3-(N,N-dicyclobutylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran hydrochloride (0.77 g, 2.26 mmol) was dissolved in anhydrous CH_2Cl_2 (20 mL) and cooled to -40°C . To the solution was BBr_3 (0.54 mL, 5.7

mmol), dissolved in anhydrous CH_2Cl_2 (3 mL), added dropwise. The cooling-bath was removed and after 2 h at room temperature the reaction was complete.

5 The reaction was poured out onto an ice/2 M NH_3 solution and the mixture was extracted, twice, with diethyl ether. The combined ether portions were dried (MgSO_4), filtered, and the solvent removed in vacuo to give the crude residue.

10 Chromatography on silica (eluent: 50% ethyl acetate/hexane) gave 0.58 g (89% yield) of the title compound as a white solid. Mp: 170-2°C.

$[\alpha]_D^{21} = -114.4^\circ$ (C=0.1; CHCl_3). GC-MS (70 eV) M= 291 (2%).

15 **c) (R)-3-(N,N-Dicyclobutylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran**

(R)-3-(N,N-Dicyclobutylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran (0.59 g, 2.02 mmol) and collidine (0.37 mL, 2.8 mmol) were dissolved in 20 anhydrous CH_2Cl_2 (40 mL) and cooled to -40°C.

Trifluoromethanesulfonic anhydride (0.41 mL, 2.4 mmol) was added dropwise and allowed to warm to ambient temperature, and after coming to 0°C the reaction was 25 done. The reaction was diluted with CH_2Cl_2 and washed with a saturated aqueous NaHCO_3 solution, dried (MgSO_4), filtered, and evaporated in vacuo to give a crude residue. Chromatography on silica (eluent: CH_2Cl_2) gave 0.84 g (99% yield) of the title compound 30 as a clear oil.

$[\alpha]_D^{21} = -90.9^\circ$ (C=0.1; CHCl_3). GC-MS (70 eV) M= 423 (3%).

35 **Methyl (R)-3-(N,N-Dicyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate**

(R)-3-(N,N-Dicyclobutylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (0.82 g, 1.94 mmol) was dissolved in a solution of

DMF/methanol (6:2, 15 mL) and then degassed followed by the inlet of carbonmonoxide (x3). With a slight positive pressure of carbonmonoxide, palladium(II)-acetate (14 mg), 1,3-bis(diphenylphosphino)propane (25 mg) and triethylamine (0.60 mL, 4.3 mmol) were added and the reaction mixture was degassed and subjected to carbonmonoxide once again. The reaction was heated to 70°C under carbonmonoxide atmosphere with vigorous stirring for 5.5 h.

The reaction was allowed to cool and the solvent was removed in vacuo. The remains were taken into a 2 M solution of NH₃ and then extracted, twice, with diethyl ether. The combined ether portions were dried (MgSO₄), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 12.5% ethyl acetate/hexane) gave 501 mg (78% yield) of the title compound as a clear oil.
 $[\alpha]_D^{21} = -138.2^\circ$ (C=0.1; CHCl₃). GC-MS (70 eV) M= 333 (4%).

e) (R)-5-Carbamoyl-3-(N,N-dicyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

Methyl (R)-3-(N,N-dicyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate (490 mg, 1.47 mmol) was refluxed with a 6 M solution of HCl (20 mL) for 3.5 h. The solution was cooled, concentrated to dryness in vacuo and anhydrous toluene was added and the solvent was removed in vacuo (x4).

To the white solid was thionyl chloride (15 mL) added and the solution was allowed to stir at room temperature overnight. The excess thionyl chloride was removed in vacuo, anhydrous toluene added and the solvent removed in vacuo.

The acid chloride was dissolved in CH₂Cl₂ (20 mL) and added dropwise to a cooled solution (ice-bath) of concentrated NH₃ (20 mL). The reaction was allowed to stir at room temperature for 30 min. The CH₂Cl₂ phase

was separated and the aqueous portion was re-extracted with CH_2Cl_2 (x3). The combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and evaporated in vacuo to give the crude residue. Chromatography on silica (eluent: ethyl acetate) gave 430 mg (92% yield) of a white solid. Mp: 141.2-142.2°C.

$[\alpha]_D^{21} = -151.5^\circ$ (C=0.1; CHCl_3). GC-MS (70 eV) M= 318 (3%).

The hydrochloride salt was made by dissolving the pure base in ether and dropping an excess of an ethereal HCl solution. The salt was washed with diethyl ether to give a white solid. Mp: 120°C sinters.

Example 5

(R)-5-Carbamoyl-3-(N-cyclobutyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

a) (R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclobutylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (1.01 g, 4.02 mmol) was dissolved in anhydrous methanol (20 mL) and to this was n-propionaldehyde (3.0 mL, 40.2 mmol) added. After 1 h the reaction was cooled (ice-bath) then sodium cyanoborohydride (0.46 g, 7.24 mmol) was added, the pH was adjusted to 4-5 with acetic acid and the reaction was allowed to stir at room temperature over the weekend. The solvent was removed in vacuo, the remains were taken into a 2 M solution of NH_3 and then extracted thrice with diethyl ether. The combined ether portions were dried (MgSO_4), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 11% ethyl acetate/hexane) gave 0.95 g (80% yield) of the title compound as a clear oil.

$[\alpha]_D^{21} = -95.4^\circ$ (C=0.1; CHCl_3). GC-MS (70 eV) M= 293 (1%).

b) (R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran hydrochloride (1.0 g, 3.03 mmol) was dissolved in anhydrous CH_2Cl_2 (25 mL) and cooled to -40°C . To the solution was BBr_3 (0.72 mL, 7.6 mmol), dissolved in anhydrous CH_2Cl_2 (4 mL), added dropwise. The cooling-bath was removed and after 2 h at room temperature the reaction was complete. The reaction was poured out onto an ice/2 M NH_3 solution and the CH_2Cl_2 portion was separated, the aqueous layer re-extracted, twice, with CH_2Cl_2 . The combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 25% ethyl acetate/hexane followed by 50% ethyl acetate/hexane) gave 0.83 g (98% yield) of the title compound as a gum. $[\alpha]_D^{21} = -80.5^\circ$ ($C=0.1$; CHCl_3). GC-MS (70 eV) $M = 279$ (0.2%).

c) (R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran (0.80 g, 2.86 mmol) and collidine (0.53 mL, 4.0 mmol) were dissolved in anhydrous CH_2Cl_2 (30 mL) and cooled to -40°C . Trifluoromethanesulfonic anhydride (0.60 mL, 3.6 mmol) was added dropwise and allowed to warm to ambient temperature, and after coming to 0°C the reaction was done. The reaction was diluted with CH_2Cl_2 and washed with a saturated aqueous NaHCO_3 solution, the aqueous was re-extracted, twice, with CH_2Cl_2 the combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and evaporated in vacuo to give a crude residue. Chromatography on silica (eluent: CH_2Cl_2) gave 1.01 g (86% yield) of the title compound as a clear oil.

$[\alpha]^{21}_D = -78.6^\circ$ (C=0.1; CHCl₃). GC-MS (70 eV) M= 411 (1%).

5 **d) Methyl (R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate**

(R)-3-(N-Cyclobutyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (1.00 g, 2.43 mmol) was dissolved in a solution of 10 DMF/methanol (6:2, 20 mL) and then degassed followed by the inlet of carbonmonoxide (x3). With a slight positive pressure of carbonmonoxide, palladium(II)-acetate (18 mg), 1,3-bis(diphenylphosphino)propane (25 mg) and triethylamine (0.75 mL, 5.3 mmol) were added 15 and the reaction mixture was degassed and subjected to carbonmonoxide once again. The reaction was heated to 70°C under carbonmonoxide atmosphere with vigorous stirring for 6 h.

The reaction was allowed to cool and the solvent was 20 removed in vacuo. The remains were taken into a 2 M solution of NH₃ and then extracted, twice, with diethyl ether. The combined ether portions were dried (MgSO₄), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 15% 25 ethyl acetate/hexane) gave 0.73 mg (94% yield) of the title compound as a clear oil.

$[\alpha]^{21}_D = -130.1^\circ$ (C=0.1; CHCl₃). GC-MS (70 eV) M= 321 (2%).

30 **e) (R)-5-Carbamoyl-3-(N-cyclobutyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran**

Methyl (R)-3-(N-cyclobutyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate (0.71 mg, 35 2.21 mmol) was refluxed with a 6 M solution of HCl (30 mL) for 3.5 h. The solution was cooled, concentrated to dryness in vacuo and anhydrous toluene was added and the solvent was removed in vacuo (x4).

To the white solid was thionyl chloride (20 mL) added

and the solution was allowed to stir at room temperature overnight. The excess thionyl chloride was removed in vacuo, anhydrous toluene added and the solvent removed in vacuo.

5 The acid chloride was dissolved in CH_2Cl_2 (30 mL) and added dropwise to a cooled solution (ice-bath) of concentrated NH_3 (30 mL). The reaction was allowed to stir at room temperature for 40 min. The CH_2Cl_2 phase was separated and the aqueous portion was re-extracted
10 with CH_2Cl_2 (x3). The combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and evaporated in vacuo to give the crude residue. Chromatography on silica (eluent: ethyl acetate) gave 622 mg (92% yield) of a white semi-crystalline solid. A portion was
15 recrystallized from ethyl acetate/hexane to give a feathery white solid. Mp: 107-9°C.
[α] $^{21}_{\text{D}}$ = -133.0° (C=0.1; CHCl_3). GC-MS (70 eV) M= 306 (0.5%).

20 Example 6

(R)-5-Carbamoyl-3-(N-cyclobutyl-N-isopropylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

25 a) (R)-8-Fluoro-3-(N-isopropylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran

(R)-3-Amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (1.62 g, 8.21 mmol) was dissolved in anhydrous methanol (20 mL) and to this was acetone (6.0
30 mL, 82.1 mmol) added. The reaction was cooled (ice-bath) then sodium cyanoborohydride (0.92 g, 14.8 mmol) was added, the pH was adjusted to 4-5 with acetic acid and the reaction was allowed to stir at room temperature overnight. The solvent was removed in
35 vacuo, the remains were taken into a 2 M solution of NH_3 and then extracted thrice with diethyl ether. The combined ether portions were dried (MgSO_4), filtered, and the solvent removed in vacuo to give the crude residue that was used as is in the next reaction.

GC-MS (70 eV) M= 239 (81%).

b) (R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

5
10
15
20
25
(R)-8-Fluoro-3-(N-isopropylamino)-5-methoxy-3,4-dihydro-2H-1-benzopyran (1.96 g, 8.19 mmol) was dissolved in anhydrous methanol (20 mL) and to this was cyclobutanone (6.1 mL, 81.9 mmol) added. The reaction was cooled (ice-bath) then sodium cyanoborohydride (2.0 g, 16.4 mmol) was added, the pH was adjusted to 4-5 with acetic acid, 3 Å molecular sieves were added and the reaction was allowed to stir at room temperature overnight. After 24 h the pH was again adjusted to 4-5 and the reaction was allowed to stir for 3 more days. The reaction was filtered, solvent was removed in vacuo, the remains were taken into a 2 M solution of NH₃ and then extracted thrice with diethyl ether. The combined ether portions were dried (MgSO₄), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 10% ethyl acetate/hexane) gave 1.60 g (77% yield) of the title compound as a clear oil.
[α]_D²¹ = -95.1° (C=0.1; CHCl₃). GC-MS (70 eV) M= 293 (3%).

c) (R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran

30
35
(R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran hydrochloride (1.76 g, 5.34 mmol) was dissolved in anhydrous CH₂Cl₂ (45 mL) and cooled to -40°C. To the solution was BBr₃ (1.3 mL, 13.4 mmol), dissolved in anhydrous CH₂Cl₂ (7 mL), added dropwise. The cooling-bath was removed and after 2 h at room temperature the reaction was complete. The reaction was poured out onto an ice/2 M NH₃ solution and the CH₂Cl₂ portion was separated, the aqueous layer re-extracted, twice, with CH₂Cl₂. The

combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and the solvent removed in vacuo to give the crude residue.

5 Chromatography on silica (eluent: 30% ethyl acetate/hexane) gave 1.46 g (98% yield) of the title compound as a gum. $[\alpha]_D^{21} = -95.7^\circ$ ($C=0.1$; CHCl_3). GC-MS (70 eV) $M = 279$ (0.7%).

10 The hydrochloride salt was made by dissolving the pure base in ether and dropping an excess of an ethereal HCl solution. The salt was washed with diethyl ether to give a white solid Mp: 120°C sinters.

15 **d) (R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran**

(R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran (1.36 g, 4.87 mmol) and collidine (0.90 mL, 6.8 mmol) were dissolved in anhydrous CH_2Cl_2 (50 mL) and cooled to -40°C .

20 Trifluoromethanesulfonic anhydride (1.05 mL, 6.1 mmol) was added dropwise and allowed to warm to ambient temperature, and after coming to 0°C the reaction was done. The reaction was diluted with CH_2Cl_2 and washed with a saturated aqueous NaHCO_3 solution, the aqueous
25 was re-extracted, twice, with CH_2Cl_2 the combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and evaporated in vacuo to give a crude residue.

30 Chromatography on silica (eluent: 70% hexane/ CH_2Cl_2) gave 1.67 g (83% yield) of the title compound as a slightly yellow oil.

$[\alpha]_D^{21} = -86.8^\circ$ ($C=0.1$; CHCl_3). GC-MS (70 eV) $M = 411$ (0.3%).

35 **e) Methyl (R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate**

(R)-3-(N-Cyclobutyl-N-isopropylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (1.65 g, 4.01 mmol) was dissolved in a solution of

DMF/methanol (6:2, 30 mL) and then degassed followed by the inlet of carbonmonoxide (x3). With a slight positive pressure of carbonmonoxide, palladium(II)-acetate (30 mg), 1,3-bis(diphenylphosphino)propane (55 mg) and triethylamine (1.25 mL, 8.8 mmol) were added and the reaction mixture was degassed and subjected to carbonmonoxide once again. The reaction was heated to 70°C under carbonmonoxide atmosphere with vigorous stirring for 6 h.

The reaction was allowed to cool and the solvent was removed in vacuo. The remains were taken into a 2 M solution of NH_3 and then extracted, twice, with diethyl ether. The combined ether portions were dried (MgSO_4), filtered, and the solvent removed in vacuo to give the crude residue. Chromatography on silica (eluent: 8% ethyl acetate/hexane) gave 1.18 mg (92% yield) of the title compound as a clear oil.
 $[\alpha]_D^{21} = -139.1^\circ$ ($C=0.1$; CHCl_3). GC-MS (70 eV) $M = 321$ (3%).

f) (R)-5-Carbamoyl-3-(N-cyclobutyl-N-isopropylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

Methyl (R)-3-(N-cyclobutyl-N-isopropylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate (1.16 mg, 3.61 mmol) was refluxed with a 6 M solution of HCl (30 mL) for 3.5 h. The solution was cooled, concentrated to dryness in vacuo and anhydrous toluene was added and the solvent was removed in vacuo (x4).

To the white gum was thionyl chloride (35 mL) added and the solution was allowed to stir at room temperature overnight. The excess thionyl chloride was removed in vacuo, anhydrous toluene added and the solvent removed in vacuo.

The acid chloride was dissolved in CH_2Cl_2 (50 mL) and added dropwise to a cooled solution (ice-bath) of concentrated NH_3 (50 mL). The reaction was allowed to stir at room temperature for 40 min. The CH_2Cl_2 phase was separated and the aqueous portion was re-extracted

with CH_2Cl_2 (x3). The combined CH_2Cl_2 portions were dried (MgSO_4), filtered, and evaporated in vacuo to give the crude residue. Chromatography on silica (eluent: ethyl acetate) gave 1.06 g (95% yield) of a white foam. The foam was crystallized using CH_2Cl_2 /hexane to give a white solid. Mp: 127.8-128.4°C. $[\alpha]_{\text{D}}^{21} = -143.0^\circ$ (C=0.1; CHCl_3). GC-MS (70 eV) M= 306 (0.3%).

The hydrochloride salt was made by dissolving the pure base in ether and dropping an excess of an ethereal HCl solution. The salt was washed with diethyl ether to give a white solid. Mp: 120°C sinters.

Example 7

(R)-5-Carbamoyl-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

a) (R)-3-(N-Cyclopentylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

To a solution of (R)-3-amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (2.5 g, 12 mmol), cyclopentanone (3.3 g, 36 mmol) and HOAc (0.7 g, 12 mmol) in methanol (25 mL) was NaCNBH_3 (2.5 g, 40 mmol) added in portions at room temperature. The solution was stirred at room temperature overnight to give a quantitative yield of the title compound. GC/MS (70 eV) M=265 (30%).

b) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

To the solution of (R)-3-(N-cyclopentylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran in methanol (25mL) were propanal (2 g, 36 mmol) and NaCNBH_3 (2 g, 40 mmol) added. The solution was stirred overnight to give the desired compound in a 97% yield according to GC. The solvent was removed in vacuo and the residue

was worked up by extraction to give 3.7 g of the title compound as a colorless oil.

GC/MS (70 eV) M= 307 (40%).

5 **c) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran**

The HCl salt of (R)-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2-H-1-benzopyran was prepared by adding an excess of an ethereal HCl into an ethereal solution of the base. The solvent was evaporated in vacuo and the residue dissolved in 48% aqueous HBr (50 mL). The solution was stirred at 120°C for 1.5 h. The solution was neutralized by carefully adding conc. ammonia. Extractive work-up gave a brown oil which was filtered through a plug of silica (ethyl acetate as the eluent). The title compound was isolated (3.7 g) as a slightly yellow oil.

GC/MS (70 eV) M= 293 (40%).

20 **d) (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran**

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran was dissolved in diethyl ether (100mL). Triethylamine (3 g, 30 mmol) was added and the mixture was cooled to -20 °C. Trifluoromethanesulfonic anhydride (4.2 g, 15 mmol), dissolved in diethyl ether (20 mL), was added dropwise (5 min). After stirring for 30 min, the dark-brownish mixture was poured into water. The organic layer was separated. Flash chromatography (ethyl acetate as the eluent) gave 3.7 g of the title compound as a yellow oil in a 69% yield. GC/MS (70 eV) M= 425 (10%).

35 **e) Methyl (R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate**

(R)-3-(N-Cyclopentyl-N-n-propylamino)-8-fluoro-5-

trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (3.7 g, 8.7 mmol), DMF (50 mL), triethylamine (2.5 g, 25 mmol), methanol (4 g, 130 mmol), palladium(II)acetate (100 mg, 0.45 mmol) and 1,3-bis-(diphenylphosphino)propane (200 mg, 0.48 mmol) were placed in a round-bottomed flask. The solution was stirred at 75°C in an atmosphere of carbon monoxide for 4 h. After evaporation of the solvent in vacuo and subjecting the crude to flash chromatography, 2.5 g (86 % yield) of the title compound as a colorless oil was isolated. GC/MS (70 eV) M= 335 (20 %).

f) (R)-5-Carbamoyl-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

Methyl (R)-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate (1.4 g, 4 mmol) was hydrolysed (6 M HCl, refluxed for 2 h) and the solvent was removed in vacuo. The crude acid was treated with SOCl₂ (room temperature for 5 min) to form the acid chloride which after removal of excess SOCl₂ in vacuo was added to conc. ammonia to form the amide. The crude product was isolated and purified by flash chromatography. The HCl salt was prepared by adding an excess of ethereal HCl into an ethereal solution of the pure base to afford the title compound (0.5 g, yield 35%) as white crystals. Mp: 85°C dec. $[\alpha]_D^{20}$ (base) = -91° (C= 0.2; CH₂Cl₂). GC/MS (70 eV) M= 320 (25%).

Example 8

(R)-5-Carbamoyl-3-(N-cyclohexyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

a) (R)-3-(N-Cyclohexylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

To a solution of (R)-3-amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (0.45 g, 2.2 mmol),

cyclohexanone (0.7 g, 7.2 mmol) and HOAc (0.14 g, 2.3 mmol) in methanol (25 mL) was NaCNBH₃ (0.5 g, 8 mmol) added in portions at room temperature. The solution was stirred at room temperature overnight to give a
5 quantitative yield of the title compound.
GC/MS (70 eV) M= 279 (30%).

b) (R)-3-(N-Cyclohexyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

10 To the solution of (R)-3-(N-cyclohexylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran in methanol (25 mL) were propanal (1.3 g, 23 mmol) and NaCNBH₃ (0.15 g, 2.3 mmol) added. The solution was stirred overnight to
15 give the desired compound in a 97% yield according to GC. The solvent was removed in vacuo and the residue was worked up by extraction to give 0.7 g of the title compound as a colorless oil.
GC/MS (70 eV) M= 321 (40%).

20

c) (R)-3-(N-Cyclohexyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran

The HCl salt of the (R)-3-(N-cyclohexyl-N-n-propylamino)-8-fluoro-5-methoxy-3,4-dihydro-2-H-1-benzopyran was prepared by adding an excess of an
25 ethereal HCl into an ethereal solution of the base. The solvent was evaporated in vacuo and the residue dissolved in 48% aqueous HBr (20 mL). The solution
30 was stirred at 120°C for 1.5 h. The solution was neutralized by carefully adding conc. ammonia. Extractive work-up gave a brown oil which was filtered through a plug of silica (ethyl acetate as the eluent).
The title compound was isolated (0.6 g) as a slightly
35 yellow oil.
GC/MS (70 eV) M= 307 (40%).

d) (R)-3-(N-Cyclohexyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclohexyl-N-n-propylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran was dissolved in diethyl ether (30 mL) and triethylamine (0.8 g, 8 mmol) was added and the mixture was cooled to -20 °C. Trifluoromethanesulfonic anhydride (0.8 g, 2.8 mmol), dissolved in diethyl ether (10 mL), was added dropwise (5 min). After stirring for 30 min, the dark-brownish mixture was poured into water. The organic layer was separated. Flash chromatography (EtOAc/hexane 1+1 as the eluent) gave 0.8 g of the title compound as a yellow oil.

GC/MS (70 eV) M= 439 (20%).

e) Methyl (R)-3-(N-Cyclohexyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate

(R)-3-(N-Cyclohexyl-N-n-propylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (0.8 g, 1.8 mmol) (4), DMF (30 mL), triethylamine (0.5 g, 5 mmol), methanol (0.8 g, 13 mmol), palladium(II)acetate (30 mg, 0.14 mmol) and 1,3-bis-(diphenylphosphino)propane (60 mg, 0.14 mmol) were placed in a round-bottomed flask. The solution was stirred at 75 °C in an atmosphere of carbon monoxide for 4 h. After evaporation of the solvent in vacuo and subjecting the crude to flash chromatography, 0.6 g (76% yield) of the title compound as a colorless oil was isolated. GC/MS (70 eV) M= 349 (30 %).

f) (R)-5-Carbamoyl-3-(N-cyclohexyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

Methyl (R)-3-(N-cyclohexyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate (5) (0.6 g, 1.7 mmol) was subjected to alkaline hydrolysis (2% KOH in EtOH, refluxed for 2 h). The solvent was removed in

vacuo and the crude acid was treated with SOCl_2 (room temperature for 5 min) to form the acid chloride which after removal of excess SOCl_2 in vacuo was added to conc. ammonia to form the amide. The crude product was isolated and purified by flash chromatography. The HCl salt was prepared by adding an excess of ethereal HCl into an ethereal solution of the pure base to afford the title compound (86 mg, yield 14%) as white crystals. Mp: 75°C dec. $[\alpha]_D^{20} = -73^\circ$ (C= 0.2, CH_2Cl_2). GC/MS (70 eV) M= 334 (35%).

Example 9

(R)-5-Carbamoyl-3-(N-cyclopentyl-N-cyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

a) (R)-3-(N-Cyclopentylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

To a solution of (R)-3-amino-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (0.7 g, 3.4 mmol), cyclopentanone (0.7 g, 8.3 mmol) and HOAc (0.2 g, 3.5 mmol) in methanol (25 mL), NaCNBH_3 (0.7 g, 10 mmol) was added in portions at room temperature. The solution was stirred at room temperature over night. The methanol was evaporated. The residue was dissolved in EtOAc and washed with water. The organic layer was dried with Na_2SO_4 and the solvent was evaporated to give 0.9 g (100% yield) of the title compound as a colourless oil. GC indicated 99.6 purity. GC/MS (70 eV) M= 265 (30%).

b) (R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran

To a solution of (R)-3-(N-cyclopentylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (0.9 g, 3.4 mmol), HOAc (0.22 g, 3.6 mmol) and cyclobutanone (2g, 30 mmol) in methanol (25 mL), were NaCNBH_3 (1 g, 16 mmol) added in portions at room temperature. After stirring for four days GC indicated 37% product. pH was adjusted to

5 (HOAc) and additional (1 g, 15 mmol) cyclobutanone was added. After stirring for further 6 days, GC indicated 64% conversion. The solvent was evaporated and the residue worked up by extraction. Flash chromatography (EtOAc/P-ether, 1+1), gave 0.53 g (53% yield) of the title compound as a colourless oil. GC/MS (70 eV) M= 319 (3%).

10 c) (R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-5-methoxy-3,4-dihydro-2H-1-benzopyran (0.53 g, 1.6 mmol) was dissolved in 47% HBr (15 mL) and stirred at 120°C for 1.5 h. The solution was cooled by adding ice and alkalisied by 14 M ammonia. Extractive work-up gave 0.5 g of the title compound as a slightly brown oil. IR contained a typical OH-band at 3654 cm⁻¹.

20 d) (R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran

(R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-5-hydroxy-3,4-dihydro-2H-1-benzopyran was dissolved in a mixture of diethyl ether and methylene chloride (90+10, 20 mL) and triethylamine (0.7 g, 7 mmol) was added and the mixture was cooled to -20 °C.

Trifluoromethanesulfonic anhydride (0.85 g, 3 mmol), dissolved in diethyl ether (10 mL), was added dropwise (5 min). After stirring for 30 min, the dark-brownish mixture was poured into water. The solvent was evaporated. The residue was dissolved in hexane and treated with active charcoal. Filtration through celite and evaporation of the solvent afforded 0.67 g of the title compound as a colorless oil. GC/MS (70 eV) M= 437 (1%).

e) Methyl (R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate

(R)-3-(N-Cyclopentyl-N-cyclobutylamino)-8-fluoro-5-trifluoromethylsulfonyloxy-3,4-dihydro-2H-1-benzopyran (0.67 g, 1.5 mmol) (4), DMF (20 mL), triethylamine (0.6 g, 6 mmol), methanol (0.8 g, 12.7 mmol), palladium(II)acetate (22 mg, 0.1 mmol) and 1,3-bis-(diphenylphosphino)propane (44 mg, 0.1 mmol) were placed in a round-bottomed flask. The solution was stirred at 75°C in an atmosphere of carbon monoxide for 4 h. The solvent was removed in vacuo, the residue was dissolved in diethyl ether and treated with active charcoal. Evaporation of the solvent afforded 380 mg of the title compound as an uncolored oil.
GC/MS (70 eV) M= 347 (3 %).

f) (R)-5-Carbamoyl-3-(N-cyclopentyl-N-cyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran

Methyl (R)-3-(N-cyclopentyl-N-cyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran-5-carboxylate (1.4 g, 4 mmol) was hydrolysed (6 M HCl, refluxed for 2 h) and the solvent was removed in vacuo. After drying in air at room temperature over night the crude amino acid hydrochloride was treated with SOCl₂ (room temperature for 5 min) to form the acid chloride which after removal of excess SOCl₂ in vacuo was dissolved in CH₂Cl₂ and added to conc. ammonia to form the amide. The crude product was isolated and purified by flash chromatography to give 220 mg of an uncolored oil which crystallized upon standing. Recrystallization from a mixture of diethyl ether and hexane gave white crystals of the title compound.
Yield: 110 mg. Mp: 138-140°C.
[α]_D²⁰ = - 146° (C= 0.2, CH₂Cl₂).

PharmacologyPharmacological treatment of depression in man

5 Evidence is available that in depressed patients the
neurotransmission in the central nervous system (CNS)
may be disturbed. These disturbances appear to involve
the neurotransmitters noradrenaline (NA) and 5-
10 hydroxytryptamine (5-HT). The drugs most frequently
used in the treatment of depression are considered to
act by improving the neurotransmission of either or
both of these physiological agonists. Available data
suggest that the enhancement of 5-HT neurotransmission
will primarily improve the depressed mood and anxiety,
15 whereas the enhancement of noradrenaline neuro-
transmission will rather improve the retardation
symptoms occurring in depressed patients. In recent
years many efforts have been made to develop new drugs
with high selectivity for improvement of 5-HT
20 neurotransmission in the CNS.

The dominating mechanism of action for the drugs
generally used today in the therapy of mental
depression is by blocking the reuptake of the
25 endogenous neurotransmitters (NA and/or 5-HT) released
from nerve terminals in the CNS, thus increasing the
concentration of these transmitters in the synaptic
cleft and hence restoring an adequate neurotrans-
mission.

30 A fundamentally different way to improve the
neurotransmission in the central 5-HT-neurons would be
to use a direct 5-HT-receptor antagonist. In order to
minimize side effects, a high selectivity for this kind
35 of receptors would then be preferable.

The receptor activity profile, even within a specific
subreceptor group such as 5-HT_{1A} is unpredictable and
may result in different pharmacological profiles. For

example, two compounds which seem to be closely related in chemical structure may act as agonists on the 5-HT_{1A} autoreceptor while one of these compounds may act as an agonist and the other acts as antagonist on the 5-HT_{1A} postsynaptic receptor. A compound which acts as an antagonist at both pre- and postsynaptic 5-HT_{1A} receptors is by definition an antagonist. Each of these compounds would have a different pharmacological profile and be useful in the treatment of different medical conditions. The differences in receptor activity profiles and resulting pharmacological profiles is set out for the tested compounds in the analysis of the data below.

(i) 5-HT_{1A} receptor binding assay

In order to evaluate the affinity for the 5-HT_{1A}-receptor the assay described below using rat brain were used and the K_i-value were measured as shown in table 1.

Tissue preparation: Cerebral cortex and hippocampus from Sprague-Dawley rats were dissected and homogenized in 15 ml ice-cold 50 mM Tris-HCl buffer, pH 7.5, containing 4.0 mM CaCl₂ and 5.7 mM ascorbic acid ("Buffer A") with an Ultra-Turrax (Janke & Kunkel, Staufen, FRG) for ten sec. After centrifugation for 12.5 min at 17,000 rpm (39,800 x g in a Beckman centrifuge with a chilled JA-17 rotor (Beckman, Palo Alto, CA, USA)), the pellets were resuspended in buffer A and homogenization and centrifugation repeated. Each pellet was suspended in 5 ml ice-cold 0.32 M sucrose and homogenized for 5 sec. The homogenized samples were kept frozen at -70°C. When used, they were diluted with buffer A to 8 mg tissue/ml and homogenized for 10 sec. The tissue homogenates were incubated for then min at 37°C and then supplied with 10 µM pargyline followed by reincubation for 10 min.

The binding assay followed that described by Peroutka, J. Neurochem. 47, 529-540, (1986), a copy of which is attached as Exhibit 2. Essentially this assay measures the ability of a given competitor molecule to inhibit the binding of ^3H -8-OH-DPAT to 5-HT_{1A} receptors. The incubation mixture (2 ml) contained ^3H -8-OH-DPAT (0.25 to 8 nM), the desired concentration of test (competitor) compound and 5 mg/ml tissue homogenate in 50 mM Tris-CHl buffer, pH 7.5, containing 4.0 mM CaCl₂ and 5.7 mM ascorbic acid. Six different concentrations of ^3H -8-OH-DPAT were analyzed. Binding experiments were started by the addition of tissue homogenate and followed by incubation at 37°C for then min. The incubation mixtures were filtered through Whatman GF/B glass filters with a Brandel Cell Harvester (Gaithersburg, MD, USA). The filters were washed twice with 5 ml ice-cold 50 mM Tris-HCl buffer, pH 7.5, and counted with 5 ml Ultima GoldTM (Packard) in a Beckman LS 3801 scintillation counter. Nonspecific binding was measured by the addition of 10 μM 5-HT to the reaction mixture. The binding data was processed by non-linear least squares computer analysis (Munson and Rodbard, Anal. Biochem. 107, 220-239, (1980)).

Data are presented as K_i values (nM) which are calculated from the IC₅₀ value with corrections made for the concentration of the ligand and its affinity constant K_D . The IC₅₀ value is the concentration of competitor/inhibitor molecule which is sufficient to bind and effectively block one-half the receptor molecules. Each K_i value for a given test compound was obtained by performing the binding assay in duplicate at 10 different concentrations.

Table 1

	<u>Examples</u>	<u>K_i-value (nM)</u>
5	1	1.76
	2	5.45
	3	1.07
	4	< 0.3
	5	1.17
10	6	1.75
	7	1.5
	8	2.53
	9	1.52

Table 1 shows that the claimed exemplified compounds have high binding affinity to the 5-HT_{1A}-receptors.

(ii) Blockade of 5-HT synthesis

The rate of synthesis of 5-hydroxytryptamine (5-HT: serotonin) and dopamine/noradrenaline (DA/NA) is measured as the accumulation of 5-hydroxytryptophan (5-HTP) and (DOPA) 3,4-dihydroxyphenylalanine, (5-HTP) respectively during 30 min after inhibition of aromatic L-amino acid decarboxylase by m-hydroxbenzylhydrazine 2HCl (100 mg/kg i.p.); purchased from Sigma. The test substance is administrated 30 min before m-hydroxbenzylhydrazine 2HCl. The regions of the brain to be examined are rapidly dissected, frozen on dry ice and stored at -70°C until assayed.

DOPA, 5-HTP and their metabolites are extracted from brain tissue with perchloric acid, containing an internal standard (Isoprenalin), The supernatant from brain homogenate is injected into a liquid chromatographic system comprising a precolumn and an analytical column. The catechol- and indolamines are detected by coulometric oxidation.

Antagonist at presynaptic 5-HT_{1A} receptors

Minimal effective dose (MED) for blockade of 8-OH-DPAT induced decrease in 5-HT synthesis rate (administered to rats subcutaneously (SC)/orally (PO)).

5

Table 2

	<u>Examples</u>	<u>SC/PO (mg/kg)</u>
	1	3/-
10	2	3/-
	3	1/3
	4	1/10
	5	1/10
	6	1/10
15	7	3/-
	8	-
	9	1/10

Table 2 shows the minimal effective dose after subcutaneous administration for obtaining a significant effect in relation to the minimal effective dose as required to obtain a significant effect after oral administration. The ratio indicates that the claimed exemplified compounds are effective antagonists at the presynaptic 5-HT_{1A}-receptors also after oral administration.

(iii) Block 8-OH-DPAT induced temperature (antagonist)

In each test, thirty rats, weighing approx 250 g, housed in 6 cages of 5 rats, are used. The rats have free access to food and water. Before the start of testing, they are numbered and left undisturbed for at least 1 hour. Before the administration of the compound, the body temperature of each rat is measured using a YSI 2100 tele-thermometer. The thermometer probe is inserted 10 cm into the rectum and left in place for thirty seconds. The drug is then administered either subcutaneously or orally. In each experiment

vehicle and 4 doses of drug are tested. One rat in each cage receives each treatment. the order of treatment is rotated since disturbance to the cage increases the activity of the rats, and thereby their body temperature. Thirth minutes after drug administration the rats' body temperature are measured again. The procedure is repeated 60, 90 and 120 minutes after drug administration. The resultant data on body temperature is subjected to analysis of variance. A significant group by time interaction is taken as an indication of drug effect. To obtain the minimum effective dose, the mean temperature for each of the drug treated groups are compared with that of the vehicle group at each time point, using Dunnett's t-test with a level of significance of $p < 0.02$. An indication of bioavailability may be obtained by calculating the ratio between the minimum effective doses following oral and subcutaneous administration.

20 Antagonist at postsynaptic 5-HT_{1A} receptors

Minimal effective dose (MED) for blockade of 8-OH-DPAT induced temperature decrease (administered to rats subcutaneously (SC))/orally (PO).

25 Table 3

	<u>Examples</u>	<u>SC/PO (mg/kg)</u>
	1	1/3
	2	1/10
30	3	0,01/1
	4	0,03/1
	5	-
	6	0,03/3
	7	0,1/1
35	8	-
	9	-

Table 3 shows the minimal effective dose after subcutaneous administration for obtaining a significant

effect in relation to the minimal effective dose as required to obtain a significant effect after oral administration. The ratio indicates that the claimed exemplified compounds is effective antagonists at the
5 postsynaptic 5-HT_{1A}-receptors also after oral administration.


The conclusion from the data in the tables above shows that the claimed compounds are 5-HT_{1A}-receptor
10 antagonists, since they show affinity to the 5-HT_{1A} receptors and act as antagonists on both the presynaptic and the postsynaptic 5-HT_{1A} receptors. Furthermore, the desired effect obtained after subcutaneous as well as
after oral administration, strongly supporting a good
15 bioavailability.

Claims

1. A compound of the general formula

5

10



(I)

wherein

R₁ is n-propyl or cyclobutyl;

R₂ is isopropyl, tertiary butyl, cyclobutyl,
15 cyclopentyl or cyclohexyl;

R₃ is hydrogen;

R₄ is hydrogen or methyl;

as (R)-enantiomer in the form of free base or
20 pharmaceutically acceptable salts thereof.

2. A compound according to claim 1 wherein R₄ is
hydrogen.

25 3. A compound (R)-3-(N-tert-butyl-N-n-propylamino)-5-carbamoyl-8-fluoro-3,4-dihydro-2H-1-benzopyran in the form of free base or pharmaceutically acceptable salts thereof.

30 4. The compound according to claim 3 in the form of free base.

5. A compound (R)-5-carbamoyl-3-(N,N-dicyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran
35 in the form of free base or pharmaceutically acceptable salts thereof.

6. The compound according to claim 4 in the form of free base.


7. A compound (R)-5-carbamoyl-3-(N-cyclobutyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran in the form of free base or pharmaceutically acceptable salts thereof.
- 5
8. The compound according to claim 7 in the form of free base.
9. A compound (R)-5-carbamoyl-3-(N-cyclobutyl-N-isopropylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran in the form of free base or pharmaceutically acceptable salts thereof.
- 10
10. The compound according to claim 9 in the form of free base.
- 15
11. A compound (R)-5-carbamoyl-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran in the form of free base or pharmaceutically acceptable salts thereof.
- 20
12. The compound according to claim 11 in the form of free base.
13. A compound (R)-5-carbamoyl-3-(N-cyclopentyl-N-cyclobutylamino)-8-fluoro-3,4-dihydro-2H-1-benzopyran in the form of free base or pharmaceutically acceptable salts thereof.
- 25
14. The compound according to claim 13 in the form of free base.
- 30
15. A pharmaceutical composition containing as active ingredient the compound of any of claims 1-14 in association with a diluent, excipient or an inert carrier.
- 35
16. A compound as defined in any of claims 1-14 for use in therapy.

17. A compound as defined in any of claims 1-14 for use as a 5-HT_{1A} antagonist.
18. A compound as defined in any of claims 1-14 for
5 use in the treatment of disorders in the central nervous system especially 5-hydroxytryptamine mediated disorders.
19. A compound as defined in claim 18 for use in the
10 treatment of depression, anxiety, anorexia, bulimia, senile dementia, migraine, obsessive-compulsive disorder, stroke, Alzheimer's disease, hypertension, gastrointestinal disorders, thermoregulatory and sexual
15 disturbances, pain or disturbances in the cardiovascular system.
20. Use of a compound defined in any of claims 1-14 for the manufacture of a medicament for treatment of disorders in the central nervous system, especially
20 5-hydroxytryptamine mediated disorders.
21. Use according to claim 20 for the manufacture of a medicament for treatment of depression, anxiety, anorexia, bulmia, senile dementia, migraine,
25 obsessive-compulsive disorder, stroke, Alzheimer's disease, hypertension, gastrointestinal disorders, thermoregulatory and sexual disturbances, pain or disturbances in the cardiovascular system.
22. A method for treatment of disorders in the central nervous system, especially 5-hydroxytryptamine mediated disorders by administering to mammals and man a compound defined in any of claims 1-14.
23. A method according to claim 22 for treatment of depression, anxiety, anorexia, bulimia, senile dementia, migraine, obsessive-compulsive disorder, stroke, Alzheimer's disease, hypertension, gastrointestinal disorders, termoregulator and sexual

disturbances, pain or disturbances in the cardiovascular system.

24. A process for the preparation of the compound as defined in claim 1-14, by

i, converting directly the compound of formula IV




15

wherein Y is a leaving group such as OSO_2CF_3 , halide by catalytic cycle using a zerovalent transition metal such as Pd or Ni and treatment with carbon monoxide followed by amination,

20

ii, converting the compound of formula IV to a compound V



25 wherein Z is Cl, Br, OH or OR_p where R_p is $\text{C}_1\text{-C}_6$ alkyl, by catalytic cycle using a zerovalent transition metal such as Pd or Ni and treatment with carbon monoxide followed by amination of the compound of formula V,

30 to give the compound of formula I, whereupon if desired is converted to a salt.

25. A process according to claim 24 wherein the amination is prepared by using ammonia or methylamine, ethylamine or isopropylamine.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/01010

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 311/58, A61K 31/35

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, A1, 9109853 (AKTIEBOLAGET ASTRA), 11 July 1991 (11.07.91) --	1-21,24-25
A	WO, A1, 9307135 (AKTIEBOLAGET ASTRA), 15 April 1993 (15.04.93) --	1-21,24-25
A	WO, A1, 9015047 (THE UPJOHN COMPANY), 13 December 1990 (13.12.90), page 91, line 33 - line 34 -- -----	1-21,24-25

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

2 January 1995

Date of mailing of the international search report

10 -02- 1995

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/01010

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 22-23
because they relate to subject matter not required to be searched by this Authority, namely:
A method for treatment of the human or animal body by therapy,
see Rule 39.1.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

26/11/94

International application No.

PCT/SE 94/01010

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 9109853	11/07/91	AU-B- 641204	16/09/93
		AU-A- 6977091	24/07/91
		CA-A- 2047237	23/06/91
		CN-A- 1052669	03/07/91
		EP-A- 0460169	11/12/91
		JP-T- 4503682	02/07/92
WO-A1- 9307135	15/04/93	AU-A- 2768492	03/05/93
		CA-A- 2118708	15/04/93
		EP-A- 0538222	21/04/93
		EP-A- 0607274	27/07/94
		FI-A,D- 941616	08/04/94
		HU-D- 9400993	00/00/00
WO-A1- 9015047	13/12/90	AU-B- 654653	17/11/94
		AU-A- 5822190	07/01/91
		CA-A,A- 2051399	01/12/90
		EP-A- 0476016	25/03/92
		JP-T- 4505618	01/10/92